Table 2.32-- PREVALENCE OF VARIOUS SUBSTANCES WITH HAWAII STUDENTS: 1987 TO 2002

[In percentages. Use of substance at least once in the student's lifetime]

Substance	1987	1989	1991	1993	1996	1998	2000	2002
Any illicit drug,								
include inhalants 1/								
6th grade	14.4	11.5	9.2	12.4	13.4	13.7	8.3	9.5
8th grade	27.1	22.3	21.6	27.3	29.6	26.3	23.3	22.0
10th grade	38.7	33.3	33.5	38.7	41.3	42.9	36.9	40.4
12th grade	50.5	41.4	39.3	42.0	47.7	50.3	48.4	49.4
Any illicit drug,	00.0		00.0	12.0	.,,,	00.0	10.1	10.1
exclude inhalants 2/								
6th grade	_	_	_	_	6.4	6.6	4.2	5.2
8th grade	_	_	_	_	23.0	20.9	18.5	18.2
10th grade	_	_	_	_	37.8	40.7	35.2	38.6
12th grade	_	_	_	_	45.9	48.9	47.8	48.5
Marijuana					10.0	10.0	17.0	10.0
6th grade	3.0	2.5	1.7	2.4	5.1	4.9	2.4	2.6
8th grade	16.5	13.1	12.3	16.7	21.5	19.2	15.9	15.9
10th grade	32.9	26.2	25.7	31.4	36.5	39.2	33.2	35.8
12th grade	46.0	36.2	34.3	37.1	44.7	47.7	45.8	46.2
Methamphetamine 3/	10.0	00.2	01.0	07.1		.,,,	10.0	10.2
6th grade	_	1.4	1.0	1.3	1.4	1.8	0.5	0.4
8th grade	_	6.1	4.3	4.9	4.4	4.6	2.3	2.0
10th grade	_	9.9	7.0	7.8	5.9	6.7	4.5	4.2
12th grade	_	11.7	8.9	8.4	7.5	7.7	5.8	5.3
Any alcohol use			0.0	0			0.0	0.0
6th grade	47.6	34.6	31.1	34.9	29.8	31.6	24.2	20.0
8th grade	64.7	57.3	55.8	57.4	54.0	52.6	49.2	42.5
10th grade	76.1	71.1	72.9	73.3	73.4	72.3	67.1	64.7
12th grade	85.9	80.5	79.8	79.2	79.7	81.2	77.2	75.4
Any tobacco use	00.0	00.0	. 0.0			0		
6th grade	-	-	-	-	24.2	22.2	12.7	10.5
8th grade	_	-	-	-	46.6	44.1	37.2	28.2
10th grade	_	_	_	-	58.4	58.0	50.5	43.2
12th grade	-	-	-	-	64.7	63.6	60.0	50.5

^{1/} Includes use of marijuana, inhalants, cocaine, methamphetamine, heroin or other opiates, sedatives or tranquilizers, hallucinogens, steroids, ectasy/MDMA, GHB, Rohypnol, or ketamine. Ectasy was first added to the survey in 1998. GHB, Rohypnol, and ketamine were first added to the survey in 2002.

Source: Hawaii State Department of Health, Alcohol and Drug Abuse Division, *The 2002 Hawaii Student Alcohol, Tobacco and Other Drug Use Study (1987-2002) Hawaii Adolescent Prevention and Treatment Needs Assessment Executive Summary (2003)*, table 1.

^{2/} Includes the use of marijuana, cocaine, methamphetamine, heroin or other opiates, sedatives or tranquilizers, hallucinogens, steroids, ecstasy/MDMA, GHB, Rohypnol, or ketamine. Ectasy was first added to the survey in 1998. GHB, Rohypnol, and ketamine were first added to the survey in 2002. 3/ Commonly known as "Ice".